arXiv:1211.3993v1 [q-bio.PE] 16 Nov 2012

Evolution of male life histories and age-dependent sexual signals under female choice

Joel J. Adamson[1]
[1] *Ecology, Evolution and Organismic Biology*
CB #3280
University of North Carolina,
Chapel Hill,
NC 27599
919-843-2320[*]

Strategic models have predicted that males could benefit from age-dependent sexual advertisement following evolution of increased lifespan. Dynamical considerations may play a crucial role in the origin of age-dependent sexual signals, despite strategic advantages in populations with established signals and preferences. I investigated the problem that rare trait-bearing males may suffer low viability due to small young-age signals, restricting the favorable conditions for age-dependent trait evolution. I also ask when age-dependence will prevail during trait evolution if males bearing age-dependent traits co-occur with males carrying age-independent traits. I used numerical simulations to analyze the evolution of an age-structured haploid population with no genetic drift. Age-dependence limits the evolution of male traits to cases of relatively weak selection against the trait, but the trait fixes at smaller sizes when age-dependent than when age-independent. When mode of expression (age-dependence versus age-independence) evolved along with the trait, age-independence prevailed over much of parameter space, although mode of expression remained polymorphic at small trait sizes under weak selection. The ubiquity of age-dependent traits in nature shows that many species' life-histories satisfy the conditions for age-dependent trait evolution. My results suggest that high adult male survival facilitates sexual selection by favoring the evolution of age-dependent sexual signals under fairly broad conditions.

Keywords: sexual selection; life history

---

[*] adamsonj@ninthfloor.org

I. INTRODUCTION

Correlations between juvenile survival and adult male attractiveness have been the focus of “good genes” theories of evolution of exaggerated sexual traits (Kokko et al., 2006). Good genes theories have typically ignored details of male life history (Kokko et al., 2002); many signal traits change over the reproductive lifespan of individuals (Bonduriansky et al., 2008), potentially producing confounding correlations between intrinsic viability and attractiveness. When traits depend on both age and intrinsic viability (“condition”), females may face a choice between older, slow-growing males and younger, equally attractive males. Who provides greater genetic benefits? Current good genes theories do not specify favorable conditions for the evolution of either male reproductive strategy. Although age itself may also be an indicator subject to selection (Brooks and Kemp, 2001), a different question asks how indicator traits evolve when they depend on age, without external clues to male age. When females cannot determine male age, and signal traits depend on age and condition, the question remains of how mate choice can produce evolutionary exaggeration of signal traits.

Selection maximizes fitness under life-histories with high adult survival through age-dependent allocation of resources (Evans et al., 2011). Recent studies suggest viewing sexual signals in a life-history context, since lifetime reproductive success incorporates selective episodes from across the lifespan of an individual (Badyaev and Qvarnström, 2002; Jacob et al., 2007). Another viewpoint suggests that sexual signals create trade-offs with other life-history traits, particularly survival (Kokko, 2001; Kokko et al., 2002). Sexual selection models could therefore account for variation in age-specific investment in sexual signaling. Signals that change predictably over the individual’s lifetime form one clear example of age-dependent signaling. Data show the importance of age through a crucial role of age-dependence in the evolution of many sexual signals, and a role for sexual selection in many age-specific traits (Bonduriansky et al., 2008). Recent investigations (Evans et al., 2011; Kivleniece et al., 2010) have furthermore found strong support for strategic age-dependent or terminal investment in sexual signaling.

Although age-dependence of signals and reproductive strategies appears common in nature, it may arise due to direct or indirect selection. Evans et al. (2011) suggest alternative hypotheses explaining age-dependent development of condition-dependent traits: (1) a signal may remain tightly correlated with overall health, leading to age-dependent decline as a result of senescence; (2) males at older age classes should have higher genetic quality from surviving more episodes of selection (Brooks and Kemp, 2001), leading to the appearance of age-dependent expression of condition-dependent signals; or (3) growth, rather than differential survival, could reveal genetic differences between individuals, leading to age-dependence in a more strictly strategic sense. Life history theory generally predicts that males at older ages should invest more in sexual advertisement due to declining opportunities to breed (Partridge and Endler, 1987). However, other life-history or strategic factors may mitigate the development of age-dependent sexual display. Certain growth patterns may promote semelparity. Another possibility is that behavioral strategies may change over the lifespan, whereas body size and morphological signal expression remain constant. Rather than developing sexual traits, males could develop strategies of coercion or sperm competition that reflect terminal investment (Bonduriansky et al., 2008; Promislow, 2003; Wedell et al., 2006).

If we restrict our discussion to sexual advertisements, do certain life-histories select for age-dependent signaling? Strategic models generally predict a correlation between adult male survival and age-dependent expression of male sexual traits. Kokko (1997) found evolutionarily stable strategies for distributing signaling effort over the reproductive lifespan under fairly broad conditions. Proulx et al. (2002) also found life-history conditions favorable to age-dependent signaling. Using dynamic programming models, Proulx et al. showed that conditions for honest trait development in iteroparous males may differ qualitatively from conditions for semelparous males. Males within an age class have expectations of future breeding opportunities correlated with condition. This leads to depressed investment by young, high-condition males, and a more rapid age-dependent trajectory of trait development for lower-condition males with shorter life-expectancy. Both of these models suggest that increased lifespan favors the evolution of age-dependent sexual signaling.

Can strategic models actually explain the prevalence of age-dependent traits in nature? Age-dependent traits present a number of difficulties for current models of evolutionary dynamics under sexual selection, despite the results of strategic models: (1) increasing expression over a lifespan with high adult mortality means that age-dependent traits may rarely occur in finite (i.e. real) populations and could be lost to drift; (2) strong preferences for old males with large traits could lead to higher rates of deleterious mutation in the population (Hansen and Price, 1995, 1999); and (3) attractiveness must compensate for the cost of the trait, since without significant trait size at young ages, rare mutant males bearing age-dependent traits may not attract mates and viability selection will eliminate the trait; and (4) Age-dependence reduces heritability of sexual signals by masking genetic differences between differently-aged males of varying genotypes, i.e. males of broadly varying genotypes can express very similar traits. Thus although age-dependent signaling of condition is evolutionarily stable under high adult male survival (Kokko, 1997, 1998; Kokko and Lindström, 1996; Proulx et al., 2002), the evolutionary origin of such traits raises questions. In particular, there is a “dynamical problem” raised by the origin of age-dependent traits by mutation. First, consider a single episode of mate choice in a population with an established age-independent signal: a mutant male who displays his trait dependent on his age will not be attractive compared to other males of similar and older ages; at best he will appear to be of low condition and not attract any females. Alternatively, consider that in a population with no established male trait, a young male with an age-dependent trait will produce a small trait that fails to balance costs with attractiveness. Survival until the age where the trait is attractive enough may be quite rare. Even if the female preference is common due to genetic drift, females may never encounter such a mutant in a real population with poor mixing. Selection against the trait overwhelms advantages due to mating success in both cases.

I address this dynamical problem by finding the selective and life-history conditions where an age-dependent trait can spread. I also address the evolution of the trait with genetic variation in *mode of expression* (age-dependent versus age-*independent*), to find the conditions where age-dependence is favored directly over age-independence. I conducted numerical population genetic simulations to find sufficient conditions for the evolution of age-dependent traits. I assumed that evolution of traits, preferences, condition and mode of expression (age-dependent versus age-independent) occurs within a large population (i.e. no genetic drift) with overlapping male generations. I further assume that females cannot directly detect a male's age, and that male trait size depends on condition as well as age. I compared allele frequency change under age-dependent trait development, as well as under three different models of age-independent trait development, and simultaneously evolving age-dependent and age-independent traits. I show that age-dependent traits evolve under qualitatively different selective conditions from age-independent traits. Age-dependent traits solve the dynamical problem by producing small traits that can withstand selection at young ages in accordance with strategic life-history theory.

II. MODEL

My model extends a classic model of female choice (Kirkpatrick, 1982) by adding age-dependence, iteroparity, overlapping generations of males, and condition-dependence. The model uses a haploid life cycle (Figure 1): new zygotes arise from meiosis and undergo viability selection followed by mating. Males that survive the first round of selection and mating proceed to another round of selection and mating, and so on. Males provide no direct benefits to females, and can mate with multiple females. Each female mates once and lives for one episode of viability selection (i.e. age-0 males supply female allele frequencies). I assume a large population size so that we can ignore the effects of genetic drift. All mortality results from



FIG. 1: Haploid life cycle for cohorts of males: haploid males emerge, grow, undergo selection, then mating, then repeated episodes of selection and mating. Meiosis then produces new zygotes following mating. Mutation in condition loci occurs before selection.

selection on the trait and condition phenotypes except for males in the terminal age class, who are removed from the simulation unconditionally (see Equation (10)).

The model genome consists of five diallelic loci: two "condition" or "intrinsic viability" loci (C), the trait locus (T), the preference locus (P) and the age-dependent mode of expression locus (F). I refer to alleles of interest, e.g. the preference or trait allele, or beneficial condition alleles, with the subscript 2, and denote their frequencies by lower-case p with a subscript corresponding to the locus. For example p_{P} represents the frequency of the choosiness allele P_2. The alleles at the condition loci are either "beneficial" or "deleterious." The number of beneficial alleles across loci adds up to an individual's condition phenotype C. Biased mutation from beneficial to deleterious ($C_2 \to C_1$) occurs in condition loci at a rate of 0.001 per individual per generation; mutation occurs in the zygote stage, before the first round of selection (see Figure 1).

Males carrying the T_1 allele do not produce the trait, regardless of condition. Males produce the trait if they have the trait allele T_2. An male aged y with condition phenotype C carrying T_2 has trait size

$$t(C,y) = b\exp(Cy). \tag{1}$$

Figure 2 shows that age-specific trait size linearly depends on the constant b ("growth coefficient").

The P locus controls mate choice behavior of females. Mate choice occurs by relative preference (as in Kirkpatrick, 1982) as a function of trait size. Females carrying P_2 encountering a male with trait size t are



FIG. 2: Male trait growth for high-condition males at two values of the growth coefficient, $b = 1.0$ and $b = 0.5$. Trait size at age 0 equals b.

$\phi(t) = 1 + \alpha t$ times more likely to mate than non-choosy females. Non-choosy females, carrying P_1, mate randomly ($\phi(t) = 1$). The mating process normalizes female mating frequency such that all females have equal mating success (see Equation 5).

The F locus controls mode of expression. Males carrying the F_1 allele show age-dependent expression, whereas carriers of the F_2 allele express the trait throughout their lives at at one of three levels in a particular simulation: (1) $t(C,0) = b$, the trait value of a 0-year-old; (2) $t(C, y_{max})$, the trait value of the oldest males in the population (still dependent on condition); or (3) $\bar{t}$, the population mean trait value. I calculated $\bar{t}$ at each iteration such that males carrying F_2 contributed to the population mean as if their traits were age-dependent:

$$\bar{t} = \frac{\sum_t t f(t)}{y_{max} + 1} \tag{2}$$

where $f(t)$ describes the frequency of males with trait value t with $y_{max}+1$ age classes. Males carrying F_2 received $\bar{t}$ as their trait value for a particular episode of mating, then I updated their values in the next iteration, based on changes in $\bar{t}$. I used one level of expression per simulation, including simulations where the male population was fixed for F_2 (an age-independent population). In a different set of simulations F_2 initially occurred at a low frequency, comparable to the frequency of the trait, so that age-independent and age-dependent expression were in competition.

Summed Gaussian functions of trait and condition describe viability for males and females:

$$w_m(C,t) = 1+\exp\left(-\frac{(\mathscr{C}-C)^2}{2\mu^2}\right)+\exp\left(-\frac{t^2}{2\nu^2}\right) \tag{3a}$$

$$w_f(C) = (1-s_p)\left(1+\exp\left(-\frac{(\mathscr{C}-C)^2}{2\mu^2}\right)\right) \tag{3b}$$

where males have $\mathscr{C}$ condition loci, μ sets the relative strength of selection for condition, ν determines the relative strength of selection against the trait, and the size of s_p reflects viability costs of female choice. Smaller values of μ and ν correspond to stronger selection (see Figure 3).

The frequency of haplotypes changes through viability selection:

$$P_i'(y) = \frac{P_i(y)W_i(y)}{W(y)} \tag{4}$$

where $P_i(y)$ and $P_i'(y)$ represent the frequency of haplotype i at age y before and after selection, respectively. $W_i(y)$ correspondingly represents the viability of haplotype i at age y and $\bar{W}(y)$ represents mean viability within age class y. The matrix $\mathbf{M}$ expresses the probability of mating between a female of genotype i and a male of genotype j:

$$\mathbf{M}_{ij} = \frac{P_i' \sum_{y=0}^{y_{max}} \phi_i\left(t_j(y)\right) P_j'(y)\pi(y)}{\sum_l \sum_{\upsilon=0}^{y_{max}} \phi_i\left(t_l(\upsilon)\right) P_l'(\upsilon)\pi(\upsilon)} \tag{5}$$

where $\pi(y)$ is the frequency of males of age y. Equation (5) expresses the frequency of matings by summing over male ages the product of female mating rate (ϕ, a function of trait size) and the frequency of the male haplotype j in the general population, i.e. adjusted for the age structure π. The subscript i on mating rate refers to choosy versus non-choosy females. The denominator adds up the age-dependent sums across all male haplotypes. The ratio of these summed terms then yields the probability that a male breeding with a female of haplotype i is of haplotype j. Multiplying by the female haplotype i frequency after selection P_i^i yields the frequency of ij pairs after mate choice.


(a) $\mu = 10, \nu = 1$

(b) $\mu = 10, \nu = 0.1$

(c) $\mu = 1, \nu = 1$

(d) $\mu = 1, \nu = 0.1$

FIG. 3: Fitness surfaces for males bearing ornaments: fitness slopes away from its maximum at $t = 0$ as the trait grows, and increases with more condition alleles. Decreasing μ decreases fitness for males with fewer than 2 condition alleles. When selection on condition weakens ($\mu = 10$, (a) and (b)) the fitness profile flattens in the C-dimension; compare this to $\mu = 1$ ((c) and (d)) where males with high-viability alleles at both condition loci have significantly higher fitness than those with 1 or none. Direct selection on the trait (ν) follows a similar profile except that $\nu \leq 1$ such that only trait-less males have significantly higher fitness than trait-bearing males in all simulations.

Summing the product of mating probabilities and recombination probabilities across **M** yields the frequency of new zygotes

$$P_i^{'}(0) = \sum_{jk} R_{jk\to i} M_{jk} \tag{6}$$

where $R_{jk\to i}$ represents the proportion of jk matings that yield genotype i after recombination (Bürger, 2000). Condition loci recombine freely ($r = 0.5$) with each other and other loci; other loci recombine at arbitrary frequencies ($0 \le r \le 0.5$; see Table I).

The relative size of the zygote class is found by the age-weighted sum of the mean fecundities of all adult age classes:

$$\pi^{'}(0) = \sum_{y=0}^{y_{max}} \bar{m}(y)\pi(y) \tag{7}$$

where $m(y)$ is the fecundity of an individual aged y. The new relative size of an adult age class is given by the mean viability of the age class:

$$\begin{aligned} \pi^{'}(y) =& \bar{W}(y)\pi(y) & (8) \\ =& \pi(y)\sum_i P_i(y)W_i(y). & (9) \end{aligned}$$

I then calculate the new age distribution by dividing by the sum of all new age class sizes (Moorad and Promislow, 2008):

$$\pi^{''}(y) = \frac{\pi^{'}(y)}{\sum_{\upsilon=0}^{y_{max}} \pi^{'}(\upsilon)}. \tag{10}$$

I calculated the initial age structure by specifying λ, the geometric rate of increase for a population in stable age distribution using an arbitrary Gaussian survivorship function centered at 0:

$$l(y) = \exp\left(\frac{-y^2}{2}\right). \tag{11}$$

The age distribution is then given by (see Charlesworth, 1994):

$$\phi(y) = \frac{\lambda^{-y} l(y)}{\sum_{\upsilon=0}^{y_{max}} \lambda^{-\upsilon} l(\upsilon)}. \tag{12}$$

All simulations started with the trait allele T_2 and preference allele P_2 at non-zero frequencies in the youngest age-class, and zero in the older age-classes. Simulations ran until: (1) the trait allele T_2 fixed; (2) the trait allele T_2 or preference allele P_2 was lost (frequency dropped below 10^{-12}); (3) the preference allele P_2 fixed; (4) the Euclidean distance between successive generations in preference and trait alelel frequencies dropped below 10^{-9}; or (5) the simulation ran for 10 million iterations.

Symbol	Meaning	Values
t	Male trait size	$0-27.30$
α	Coefficient of preference	0.6,1
s_p	Coefficient of selection against choosiness	$0.0-0.0001$
μ	Coefficient of selection on condition	1,10
ν	Coefficient of selection on trait	$0.0001-1.0$ in steps of 0.02
r	Recombination fraction	$0.25, 0.5$
λ	Geometric rate of increase (initial age structure)	1.0
b	Coefficient of trait growth	$0.0-0.5$ in steps of 0.02
$\mathscr{C}$	Number of condition loci	2
y_{max}	Oldest age for males	2

TABLE I: List of variables and parameters with typical values

III. RESULTS

A. Direct selection on preference

Above $s_p = 0.0001$ the preference allele (and thereafter the trait allele) was lost at all parameter values. The behavior of the system does not appear to depend on mode of trait expression (age-dependent vs. age-independent) with $0 < s_p \leq 0.0001$, and hence for all further results $s_p = 0$ (no selection on choice).

B. Selection on male trait and growth

The first set of simulations sought to determine the important parameters for evolution of the age-dependent trait compared with an age-independent trait. The age-independent trait was the same size as the eventual size of the age-dependent trait. Age-dependent simulations ran with coefficient of preference (α), recombination frequency (r) and selection on condition (μ) at the values indicated in Table I. I used initial values of $p_C = 0.01$, $p_\mathrm{P} = 0.1$, $p_\mathrm{T} = 0.001$, and $p_\mathrm{F} = 0.0$. I compare this to a population where all initial values were the same except that males were fixed for F_2, i.e. male trait expression was age-independent and expressed at the largest size attainable ($t = b\exp(Cy_{max})$). I analyzed the relative roles of selection intensity and trait size by plotting the equilibrium value (fixation versus loss) of p_T over a plane defined by



FIG. 4: Region of fixation (dark gray) of age-dependent (a) and age-independent traits (b) in a plane defined by strength of selection against the trait (ν) and the growth coefficient of the trait (b). Dashed line indicates region of fixation under $\alpha = 0.6$; solid lines indicates region of fixation for $\alpha = 1.0$. The region of fixation for $\alpha = 0.6$ is contained within the region of fixation for $\alpha = 1.0$ in both panels. Under age-dependence b is the value of the trait for a 0-year old male. Smaller values of ν correspond to stronger selection.

$0.00001 \leq \nu \leq 1.0$ and $0 \leq b \leq 0.5$ (see Table I; see Figures 4 and 5).

The area of parameter space where the trait fixes depends on three parameters: α (strength of preference), b (size or "growth coefficient") and ν (strength of selection; Figure 4). Recombination frequency (r) and selection on condition (μ) do not appear to qualitatively affect the results in these simulations. Strength of preference affects the "readiness to mate" of females over the range of trait values present in these simulations: when $\alpha = 0.6$ a choosy female is 9.20 times more likely to mate with a 2 year-old, high-condition male with a growth coefficient of $b = 0.25$. At the higher value of $\alpha = 1.0$, a choosy female is 14.65 times more likely to mate with the same trait-bearing male. We see a notable difference between populations with age-dependent traits and populations with age-independent traits at both values of α. At $\alpha = 0.6$ the trait fixes in a very small portion of the $b - \nu$ plane near $\nu = 1.0$ in age-dependent simulations, whereas it fixes regardless of ν above the threshold b value of 0.20 in age-independent simulations. At $\alpha = 1.0$ the regions of fixation are larger, and the qualitative differences between age-dependent and age-independent expression remain: the age-dependent trait fixes in a roughly triangular region characterized by relatively weak selection and containing a region of small initial trait sizes (Figure 4a). This contrasts to age-independent simulations, where again above a threshold size ($b = 0.12$) the trait fixes independently of selection intensity (Figure 4b).

A sample of linkage disequilibrium trajectories from the four regions defined by fixation and loss, age-dependence and age-independence, shows that the trait declines until condition reaches mutation-selection balance. The male life history equilibrates by generation 30 in all cases and any variation in males is due only to mutation at the condition loci. At this point the age-dependent trait either increases to fixation or declines to extinction depending on the intensity of selection. At roughly the same point in the trajectory the age-independent trait increases or declines depending on the size of the trait.

C. Mode of expression

Another set of simulations sought to determine the crucial parameters favoring age-dependent expression over age-independent expression. Fixation of F_2 depended on the associated level of trait expression with initial polymorphism variance at the F locus (Figure 5). I simulated these conditions with initial values of $p_C = 0.01$, $p_\mathrm{P} = 0.1$, $p_\mathrm{T} = 0.1$, and $p_\mathrm{F} = 0.1$ in all age classes. Age-independent males occurred with one particular trait function ($t = b$, $t = \overline{t}$, or $t = b\exp(Cy_{max})$) for a specific simulation: when the F_2 allele caused males to have smaller average traits than males with age-dependent traits ($t(C,y) = b$ or $t(C,y) = \overline{t}$) then F_2 was lost everywhere in the b-v-plane.

When males carrying F_2 expressed the trait range of the oldest age-dependent males throughout their lives ($t = b\exp(Cy_{max})$ the F_2 allele fixed in an area including the highest intensity of selection. However, the F_2 allele also failed to increase in an area of small initial trait values where the trait did fix. In other words, the age-dependent trait fixed in an area of small trait values despite the presence of age-independent expression in the population (see Figure 5a). This effect is even stronger when selection on condition is relaxed ($\mu = 10$; see Figure 5b). The area of trait fixation where F_2 failed to increase is larger when $\mu = 10$ and includes regions of more intense selection against the trait as well as requiring larger trait sizes for fixation of F_2. Net selection seems to favor age-dependence in the areas where the young-age trait of age-dependent males is small relative to that of age-independent males, and more so when there is more variation in condition (at $\mu = 10.0$).



FIG. 5: Regions of fixation for the trait only (light gray) and the trait along with age-independent mode of expression allele (F_2; dark gray) at strong selection on condition (a) and weak selection on condition (b). At small trait sizes the trait fixes in a region where F_2 does not increase.

IV. DISCUSSION

Although age-dependent sexual traits are common in nature, their evolution presents several dynamical difficulties. Balancing the benefits derived through mate choice with the costs of growing a trait requires undergoing a period when age-dependent traits may be easily lost from the population. I have shown that an age-dependent trait can evolve under relatively weaker selection at a wider range of trait sizes than can an age-independent trait. While an age-dependent trait can evolve by producing small traits at young ages when selection is intense, larger initial trait sizes can only occur under proportionally less intense selection. This contrasts strongly with the age-independent simulations, where size of the trait primarily determines the eventual evolution of the trait. Simulations with genetic variation for mode of expression (age-dependent vs. age-independent) show that age-independence only evolves at larger trait sizes and age-dependence can predominate under weaker selection at smaller trait sizes.

Altogether the simulations show that the dynamical problem of reduced heritability of traits is “solved” by producing small traits in young males under intense selection pressure. The results suggest three factors favor the evolution of age-dependent sexual signals: (1) low adult male mortality; (2) weak selection against the ornament; and (3) strong age-dependence, i.e. small initial trait sizes. This result makes sense from a life-history strategy perspective: when adult mortality is lower, males will invest more in reproductive strategies, with a negative correlation between expected future reproduction and age-specific investment. Males can afford to invest in “frivolous” mate attraction when overall adult mortality is low. Age-dependence therefore partitions the life-history of males into two stages dominated by different fitness components: viability is most crucial at young ages and mating success is more important only later in life. This result parallels recent work on a large mammal showing that after juvenile survival, mating success during the later reproductive phase forms the most crucial fitness component (Courtiol et al., 2012). Females can benefit from choice by producing male offspring with an effectively conservative life history that allows them to survive well to old age when selection is less intense and larger, more attractive traits lead to high mating success. The equilibration of male life history is crucial. Trajectories show that the critical point in the evolution of the trait allele was when condition reached mutation-selection balance. At this point we can consider the trait an honest indicator of condition: any male carrying a deleterious mutation will carry a smaller trait and be strongly selected against in the mating phase, with the sexual selection differential on condition strongly increasing with age. As males age, they reveal additive variance in condition, thus making them more reliable signallers (Proulx et al., 2002).

The results here support the strategic modeling literature of age-dependent signals. Proulx et al. (2002) modeled the situation where male longevity and reproductive opportunities increase — e.g. under a low adult mortality environment — and younger males downplay their signaling, preserving resources for survival. Kokko (1997) came to the similar conclusion that young males of lower condition should signal more, thus obscuring the observed relationship between genetic quality and trait value. Higher-condition males can afford to downplay signaling more than low-condition males; when males signal in proportion to number of remaining reproductive attempts (a predictor of condition in any age-class), selection favors females that prefer to mate with older males. Proulx et al. allowed these strategies to compete, whereas my model uses a condition-dependent and age-dependent trait function to model the same variation. My results show that when selection weakens enough, with a particular developmental trajectory, age-dependent signaling and female preferences evolve in a population genetic model. The results showing evolution of age-dependent traits at mutation-selection balance in condition support Proulx et al.’s conclusion that under an established regime of sexual signaling, selection on life-histories will produce age-dependent signaling.

Empirical evidence and my results suggest that extending the male lifespan facilitates sexual selection. When traits are age-dependent they can develop their most exaggerated forms at older ages when selection is less intense. Age-dependent traits or mating success occur in a wide variety of taxa, including mammals (Clinton and Le Boeuf, 1993; Pemberton et al., 2004; Poissant et al., 2008), birds (Ballentine, 2009; Evans, 1997; Evans et al., 2011; Garamszegi et al., 2007; Hawkins et al., 2012; Taff et al., 2011), fish (Candolin, 2000*a*,*b*; Jacob et al., 2007; Johnson and Hixon, 2011; Miller and Brooks, 2005) and insects (Jones et al., 2000; Jones and Elgar, 2004; Judge, 2011; Kivleniece et al., 2010; Verburgt et al., 2011). Despite the demographic problems of age-dependence, widespread occurrence of age-dependent traits suggests that life-histories promoting age-dependence are common. Body size or traits directly correlated with body size (e.g. weapons and ornaments) form the most obvious example of age-dependent traits, and should satisfy the assumptions of my model. Researchers found age-dependent sexual selection based on body size in Rocky Mountain Bighorn Sheep (*Ovis canadensis*; Coltman et al. 2002), who show a typical life-history characterized by weakening viability selection and increasing heritability over the lifespan. Older males pay less of a survival cost for larger bodies and larger horn sizes, facilitating greater success in mating competition. Certain behavioral and social traits should display age-dependence under weak natural selection, such as social network connectivity (McDonald and Potts, 1994) and song repertoire (Gil et al., 2001). Age-based honesty also creates an effective constraint producing age-dependence. High-condition males cannot bypass age-dependence by “faking” the trait. Certain “skills” such as nest-building Evans (1997) may show this form of honesty.

Readers should consider some limitations of the model I used here. My model inadequately portrays situations where direct costs of female choice impact sexual selection. Direct costs of choice may considerably impact female survival and therefore a full analysis of the life-history implications of sexual selection needs to include a model of the female life-history. Indirect costs, such as increasing frequency of germ-line mutations with male age may also greatly affect female choice evolution in long-lived organisms (Beck and Promislow, 2007; Hansen and Price, 1999). Female choice may also depend on female condition, which could produce strong linkage disequilibrium between condition loci and preference loci under selection against the trait. Condition-dependent female choice could therefore broaden the range of parameters where an age-dependent trait can evolve. The trait function I used here also has somewhat narrow applicability, as it strongly favors old males: age-dependent traits in nature probably favor intermediate-aged males, who have become attractive but have not deteriorated considerably (Brooks and Kemp, 2001). A more physiologically realistic trait function (Johnson and Hixon, 2011) would peak in middle-age, but would tell us little more about the evolution of the preference without considering costs of mating with old (versus intermediate-aged) males. My model also uses an effectively infinite population size and may therefore overestimate the effect of old males in the population. The effective population size of males with honest indicators of condition may dwindle due to demographic effects in a finite population, and their alleles could easily be lost to drift.

Age-dependent signaling offers a testable hypothesis relating life-histories and sexual selection. Independent developments of iteroparity and reduced adult mortality may be associated with evolution of age-dependent sexual signals. The weakening of selection associated with lifespan development should facilitate sexual selection by concomitant reductions in selection against outrageous traits at older ages. Further modeling should ask whether senescence and accumulation of mutations could weaken this prediction.

V. ACKNOWLEDGMENTS

I would like to thank: Maria Servedio for overseeing my research and getting me to come up with an actual research question instead of just learning all the math there is to learn; Joel Kingsolver, Troy Day, Karin Pfennig, and Haven Wiley for scrutinizing my results; Sumit Dhole, Alicia Frame, Caitlin Stern, Justin Yeh, Artur Romanchuk and Daniel Promislow for providing helpful discussion and comments; Richard M. Stallman and the GNU Project for guidance with software; the developers of Guile, Trac, GNUPLOT and the GNU C Compiler for creating reliable freedom-respecting software. This research was supported by NSF DEB-0614166 and NSF DEB-0919018 to Maria Servedio.

REFERENCES

Badyaev, A. V., and A. Qvarnström. 2002. Putting sexual traits into the context of an organism: A life-history perspective in studies of sexual selection. Auk 119:301–310.

Ballentine, B. 2009. The ability to perform physically challenging songs predicts age and size in male swamp sparrows, *Melospiza georgiana*. Animal Behaviour 77:973 – 978.

Beck, C. W., and D. E. L. Promislow. 2007. Evolution of female preference for younger males. PLoS ONE 2:e939.

Bonduriansky, R., A. Maklakov, F. Zajitschek, and R. Brooks. 2008. Sexual selection, sexual conflict and the evolution of ageing and life span. Functional Ecology 22:443–453.

Brooks, R., and D. J. Kemp. 2001. Can older males deliver the good genes? Trends in Ecology & Evolution 16:308–313.

Bürger, R. 2000. The Mathematical Theory of Selection, Recombination and Mutation. J. Wiley and Sons.

Candolin, U. 2000*a*. Changes in expression and honesty of sexual signalling over the reproductive lifetime of sticklebacks. Proceedings of the Royal Society of London. Series B: Biological Sciences 267:2425 –2430.

———. 2000*b*. Increased signalling effort when survival prospects decrease: male-male competition ensures honesty. Animal Behaviour 60:417–422.

Charlesworth, B. 1994. Evolution in age-structured populations. 2nd ed. Cambridge University Press.

Clinton, W. L., and B. J. Le Boeuf. 1993. Sexual selection's effects on male life history and the pattern of male mortality. Ecology 74:1884–1892.

Coltman, D., M. Festa-Bianchet, J. Jorgenson, and C. Strobeck. 2002. Age-dependent sexual selection in bighorn rams. Proceedings of the Royal Society of London. Series B: Biological Sciences 269:165–172.

Courtiol, A., J. E. Pettay, M. Jokela, A. Rotkirch, and V. Lummaa. 2012. Natural and sexual selection in a monogamous historical human population. Proceedings of the National Academy of Sciences page Published online before print April 2012.

Evans, M. R. 1997. Nest building signals male condition rather than age in wrens. Animal Behaviour 53:749 – 755.

Evans, S. R., L. Gustafsson, and B. C. Sheldon. 2011. Divergent patterns of age-dependence in ornamental and reproductive traits in the collared flycatcher. Evolution 65:1623–1636.

Garamszegi, L. Z., J. Trk, G. Hegyi, E. Szllsi, B. Rosivall, and M. Eens. 2007. Age-dependent expression of song in the collared flycatcher, *Ficedula albicollis*. Ethology 113:246–256.

Gil, D., J. L. S. Cobb, and P. J. B. Slater. 2001. Song characteristics are age dependent in the willow warbler, *Phylloscopus trochilus*. Animal Behaviour 62:689 – 694.

Hansen, T., and D. Price. 1995. Good genes and old age: Do old mates provide superior genes? Journal of Evolutionary Biology 8:759–778.

———. 1999. Age-and sex-distribution of the mutation load. Genetica 106:251–262.

Hawkins, G. L., G. E. Hill, and A. Mercadante. 2012. Delayed plumage maturation and delayed reproductive investment in birds. Biological Reviews 87:257274.

Jacob, A., S. Nussle, A. Britschgi, G. Evanno, R. Muller, and C. Wedekind. 2007. Male dominance linked to size and age, but not to 'good genes' in brown trout *(Salmo trutta)*. BMC Evolutionary Biology 7:207.

Johnson, D. W., and M. A. Hixon. 2011. Sexual and lifetime selection on body size in a marine fish: the importance of lifehistory tradeoffs. Journal of Evolutionary Biology 24:1653–1663.

Jones, T. M., A. Balmford, and R. J. Quinnell. 2000. Adaptive female choice for middle-aged mates in a lekking sandfly. Proceedings of the Royal Society of London. Series B: Biological Sciences 267:681–686. 10.1098/rspb.2000.1056.

Jones, T. M., and M. A. Elgar. 2004. The role of male age, sperm age and mating history on fecundity and fertilization success in the hide beetle. Proceedings of the Royal Society of London. Series B: Biological Sciences 271:1311–1318.

Judge, K. A. 2011. Do male field crickets, *Gryllus pennsylvanicus*, signal their age? Animal Behaviour 81:185 – 194.

Kirkpatrick, M. 1982. Sexual selection and the evolution of female choice. Evolution 36:1–12.

Kivleniece, I., I. Krams, J. Daukste, T. Krama, and M. J. Rantala. 2010. Sexual attractiveness of immune-challenged male mealworm beetles suggests terminal investment in reproduction. Animal Behaviour 80:1015–1021.

Kokko, H. 1997. Evolutionarily stable strategies of age-dependent sexual advertisement. Behavioral Ecology and Sociobiology 41:99–107.

———. 1998. Good genes, old age and life-history trade-offs. Evolutionary Ecology 12:739–750.

———. 2001. Fisherian and good genes benefits of mate choice: how (not) to distinguish between them. Ecology Letters 4:322–326.

Kokko, H., R. Brooks, J. M. McNamara, and A. I. Houston. 2002. The sexual selection continuum. Proceedings of the Royal Society of London. Series B: Biological Sciences 269:1331–1340.

Kokko, H., M. D. Jennions, and R. Brooks. 2006. Unifying and testing models of sexual selection. Annual Review of Ecology, Evolution and Systematics 37:43–66.

Kokko, H., and J. Lindström. 1996. Evolution of female preference for old mates. Proceedings of the Royal Society of London. Series B: Biological Sciences 263:1533–1538.

McDonald, D., and W. Potts. 1994. Cooperative display and relatedness among males in a lek-mating bird. Science 266:1030–1032.

Miller, L. K., and R. Brooks. 2005. The effects of genotype, age, and social environment on male ornamentation, mating behavior, and attractiveness. Evolution 59:2414–2425.

Moorad, J. A., and D. E. L. Promislow. 2008. A theory of age-dependent mutation and senescence. Genetics 179:2061–2073.

Partridge, L., and J. Endler. 1987. Life history constraints on sexual selection. Pages 265–277 *in* Sexual selection: testing the alternatives. John Wiley and Sons Ltd.

Pemberton, J., D. Coltman, J. Smith, and D. Bancroft. 2004. Mating patterns and male breeding success. Chap. 6, pages 166–189 *in* T. Clutton-Brock and J. Pemberton, eds. Soay Sheep: Dynamics and selection in an island population. Cambridge University Press, Cambridge, UK.

Poissant, J., A. J. Wilson, M. Festa-Bianchet, J. T. Hogg, and D. W. Coltman. 2008. Quantitative genetics and sex-specific selection on sexually dimorphic traits in bighorn sheep. Proceedings of the Royal Society of London. Series B: Biological Sciences 275:623–628.

Promislow, D. 2003. Mate choice, sexual conflict, and evolution of senescence. Behavior Genetics 33:191–201.

Proulx, S., T. Day, and L. Rowe. 2002. Older males signal more reliably. Proceedings of the Royal Society of London. Series B: Biological Sciences 269:2291–2299.

Taff, C. C., C. R. Freeman-Gallant, P. O. Dunn, and L. A. Whittingham. 2011. Relationship between brood sex ratio and male ornaments depends on male age in a warbler. Animal Behaviour 81:619–625.

Verburgt, L., M. Ferreira, and J. Ferguson. 2011. Male field cricket song reflects age, allowing females to prefer young males. Animal Behaviour 81:19–29.

Wedell, N., C. Kvarnemo, C. . Lessells, and T. Tregenza. 2006. Sexual conflict and life histories. Animal Behaviour 71:999–1011.

Appendix A: Genotypic Trajectories


(a) $b = 0.1, v = 0.2$

(b) $b = 0.1, v = 0.8$


(c) $b = 0.2, v = 0.8$

(d) $b = 0.2, v = 0.2$

FIG. 6: Trajectories of linkage disequilibrium between the trait and one of the condition loci, with $\alpha = 1.0$. This quantity reflects the frequency of the genotype bearing both the trait allele and the beneficial condition allele. The four panels are defined by regions of Figure 4 bearing different patterns of fixation: (a) no fixation under age-dependence or age-independence; (b) fixation under age-dependence, not under age-independence; (c) fixation under both modes of expression; (d) fixation only under age-independence.